


## After Hip Replacement



**TRUSTED FOR OUR EXPERTISE. CHOSEN FOR OUR CARE.**



Zac Burrow, MD, DPT

I want you to know that I understand this is a big decision, and it often comes after months or even years of pain, limitation, and frustration.

**My goal is simple: to help you feel better, move better, and get back to the life you enjoy.**

I care deeply about your outcome—not just in the operating room, but in your recovery and long-term success. Every decision we make is centered around what is safest, most effective, and best for *you* as an individual.

This guide is designed to walk you through the process, answer common questions, and help you feel confident and prepared every step of the way. You are not going through this alone—we are a team, and I am committed to supporting you throughout your entire journey.

**Thank you for trusting me with your care!**

\*Pictures in this guide were obtained from AAOS, HEP2Go.com, or generated with ChatGPT\*

## General information can also be found at:

- Total Hip Replacement - OrthoInfo – AAOS  
<https://orthoinfo.aaos.org/en/treatment/total-hip-replacement>
- Total Hip Replacement - AAHKS  
<https://hipkneeinfo.org/hip-care/total-hip-replacement/>
- Video - Total Hip Replacement Animation - OrthoInfo - AAOS  
<https://orthoinfo.aaos.org/en/treatment/total-hip-replacement-animation/>

## Important Phone Numbers

- **Dr. Burrow Direct Line:** 918-927-3258
  - Call for refills of medications, general questions/concerns
  - May also use online patient portal
    - MyMobility App is not monitored
- **Surgery Scheduling:** 918-927-3301
  - Scheduler will call you to schedule surgery once all clearances are obtained
    - Call for any surgical scheduling questions
    - Cancel, reschedule, etc.
- **Billing Department:** 918-927-3737
  - Any billing questions for AOOK
- **Brace Shop:** 918-927-3308
- **Hospital Pre-Admission Testing:**
  - Oklahoma Surgical Hospital: 918-477-5073
  - St. John Broken Arrow: 918-994-8045
- **Urgent Surgical Problem:** 918-494-2665
- **Emergency:** 911

## Immediately After Surgery (Hospital Stay or Outpatient)

### Common After Surgery

- Pain and swelling in the operated leg
- A bandage covering your hip incision
- Medications to help with: Pain control & Blood clot prevention

### Physical Therapy Starts Early

- Begin **basic exercises within 24 hours**
- Help you **stand and start walking**
- Teach you how to move safely using a **walker**

### Hospital Stay

- Most patients stay **1 night in the hospital**
  - Some patients may: Go home the **same day** or may need a **longer hospital stay**

## At Home: First Few Weeks After Surgery

### Mobility & Walking

- Use a **walker at first**, then transition to a **cane** as you improve
- Walk **short distances multiple times per day**
- Gradually work toward: Longer walks & Climbing stairs safely

### Physical Therapy

- We will try to arrange **home health physical therapy**
  - This typically starts the 4<sup>th</sup> day after you return home
  - *Note: Insurance may limit this option*
- Continue:
  - **Daily home exercises**
  - Use of the **MyMobility App** for guidance

### Pain & Swelling

- It is normal to have: Pain (managed with medications and ice) & bruising / swelling
- Swelling will **improve gradually over several weeks**

### Help at Home

- You will likely need help with: Cooking, Bathing, Daily activities
- Plan for assistance for the **first 1–2 weeks**

### Activity Goals

- Walk several times per day
- Increase activity **slowly and safely**

## Mid-Term Recovery: 6 Weeks to 3 Months

### Mobility & Walking

- Walking becomes **easier and more natural**
- Decreased need for a **walker or cane**

### Return to Daily Activities

- Gradual return to normal daily routines
- May begin: Light household tasks
- Driving (*typically around 4–6 weeks, if cleared by your physician*)

### What Is Normal

- Mild **stiffness**, Some **swelling**, **Fatigue**, especially after activity
- These symptoms are common and should improve over time

### Activity Progression

- Continue to increase activity **gradually**
- Stay consistent with exercises and therapy

## Full Recovery: 3 to 12 Months

### What to Expect

- Most patients return to **normal daily activities by 3 months**
- Continued improvement in: Strength, Balance, Comfort
- Full recovery may take up to **1 year**

### Return to Activities

- Most patients can safely return to:
  - Walking
  - Swimming
  - Biking
  - Other **low-impact activities**

### Activities to Avoid

- Running
- High-impact sports
- Activities that place excessive stress on the joint

## Nutrition Before and After Surgery

Good nutrition helps your body heal, reduces complications, and improves recovery.

### General Guidelines

- If you are on a **specific diet**, continue following it
- Focus on **whole, healthy foods**
- Avoid highly processed foods (fast food, boxed, canned food, etc.)

### Recommended Nutrition Timeline

- Start **2 weeks before surgery**
- Continue for **at least 6 weeks after surgery**

### Hydration

- Avoid **alcohol and soft drinks**
- Drink **water or tea**
- Goal: **5 servings of 12 oz per day**

### Recommended Foods

- **Berries** (1–3 servings/day)
  - Blueberries, Blackberries
  - Frozen options are okay
- **Fruits** (1–2 servings/day)
  - Apples, Avocados
- **Beans** (1-2 servings/day)
  - Green beans
  - Pinto beans
- **Whole Grains** (3 servings/day)
  - Brown rice, Oatmeal
- **Vegetables** (1–2 servings/day)
  - Broccoli, Sweet potatoes
- **Greens** (2 servings/day)
  - Spinach, Kale
- **Nuts** (1 serving/day)
  - Walnuts, Pecans



### Protein (Very Important for Healing)

- Lean meats:
  - Grilled chicken / fish
- Whey or soy protein supplements

## Post-Operative Medication Instructions

These medications help control pain, prevent complications, and support healing after surgery

\*\*\*\*These are general medications. Some patients may require alternative medications or dosage\*\*\*\*

### Pain Control

- **Tylenol (Acetaminophen)** - Take **500 mg tablets – 2 tablets every 8 hours**
  - Take this **on a regular schedule**, not just when needed
- **Celebrex (Celecoxib)** - Take **200 mg once daily**
  - **Alternative (if sulfa allergy): Meloxicam 15 mg once daily**
  - If you are already taking an NSAID medication, we will just resume your current medication
  - **Important: Do NOT take Celebrex or Meloxicam if you have kidney (renal) problems**
- **Journavx (Suzetrigine)** - Non-Opioid pain medication
  - Take 1 dose of 100mg on day 1
  - Take 50mg 2 times per day after first dose
- **Gabapentin** – Take **100mg tablet three times per day**. *\*can cause drowsiness\**

### Breakthrough Pain Medication

- **Oxycodone 5 mg** - Take **1 tablet every 4 hours as needed for pain**
  - **If pain is still severe (6–10) after 30 minutes:** You may take **1 additional tablet (5 mg)**
- **Important: Do NOT take more medication until at least 4 hours after your last dose**
  - \*\*\*\*This is a **controlled substance**, and refills are limited by law\*\*\*\*
  - If a refill is needed, please call our office the day before you run out.
  - If you will need more over the weekend, call Friday morning.
    - **Do not wait until late Friday or over the weekend to ask for a refill.**

### Blood Clot Prevention

- **Aspirin 81 mg:** Take **twice daily for 28 days** and then **once daily for 14 days**. 42 total days
- **Note:** You may be given a different medication if you are high risk or already on a blood thinner

### Stomach & Side Effect Support

- **Colace (Docusate) 100 mg** - Take **1 tablet twice daily**
  - Continue while taking narcotic pain medication. Helps prevent constipation
- **Zofran (Ondansetron) 4 mg ODT** - Dissolve **1 tablet under your tongue every 8 hours as needed**
  - Use for nausea or vomiting

### Antibiotic

- **Cefadroxil 500 mg** - Take **twice daily for 7 days**
- *This may not be prescribed for all patients*

### Recommended Supplements (Over-the-Counter)

- **Vitamin D3 + K2:** 5,000 IU daily
- **Calcium Carbonate:** 1,200 mg daily

### Resume Your Regular Medications

- Restart your normal medications after surgery - **Unless your doctor tells you otherwise**



## Post-Operative Wound Care

Proper wound care is important to prevent infection and ensure good healing.

### Dressing Care

- Leave your dressing **in place until your follow-up visit (approx. 3 weeks after surgery)**
  - Unless changed by home health nursing
- **No additional wound care is needed**

### When to Call the Clinic

Call our office if you notice:

- Drainage **larger than the size of a quarter**
- Fluid **pooling under the dressing**
- Increasing redness around the incision
- **Cloudy or pus-like drainage**



### Showering

- You **may shower with your bandage on** (it is waterproof) unless told otherwise by Dr. Burrow
  - If it looks like the bandage is coming off at the edge, do **NOT** let the wound/bandage get wet
- Occasionally an incisional wound vac dressing is used. This stays on for 1 week & should **NOT** get wet

### Do NOT Submerge the Wound

- Avoid baths, Hot tubs, Swimming pools, etc. for at least **6 weeks after surgery**

### Compression Stockings

- Wear compression stockings for **6 weeks**
- Helps reduce swelling and prevent blood clots

### Skin & Incision Care

- Do **NOT** apply lotions, creams, or ointments to your incision for the first **6 weeks**

### After the Incision Heals

- Once fully healed, you may use:
  - Any scar creams or vitamin E.
- Use **sunscreen over the incision** to help prevent darkening of the scar
  - *Only after the incision is fully healed*

## Ice and Heat After Surgery

Using ice and heat correctly can help reduce pain and swelling during recovery.



### Ice (Recommended Early After Surgery)

- **How to use:**
  - Place a **towel or cloth between the ice and your skin**
  - Apply ice for **15–20 minutes at a time**
- **How often:**
  - At least **3–4 times per day**
  - You may use more often if needed
  - Early after surgery:
    - Ice every **1-2 hours** is recommended

### Heat (Avoid Early On)

- **Do NOT use heat** in the early post-operative period

### Avoid Heat Products Early

- Do NOT use: Ben-Gay, Icy Hot, Etc.

### Ice Machine (Optional)

- Ice machines can make icing easier and more consistent
  - Can add compression to improve swelling and pain
- Available for purchase at the brace shop
  - Insurance does not cover ice machines

## Activity and Exercise After Surgery

Staying active is important for your recovery, but it's important to do so safely.

*\*exercise pictures and instructions were chosen from HEP2go.com*

### Weight Bearing

- You may **put weight on your leg as tolerated**
  - Unless instructed otherwise by Dr. Burrow
- Use your comfort level as a guide

### Walking

- Walk around your home **multiple times per day**
- Short, frequent walks are best
  - This helps: Improve circulation, Reduce swelling, & Prevent blood clots

### Assistive Devices

- Use a **walker or crutches** as directed by physical therapist
  - Continue using them until cleared by physical therapy or surgical team

### Driving

- Do **NOT drive while taking pain medication**
- Do **NOT drive until cleared** by your physical therapist or physician
- **Note:** Recovery time may be **longer for right hip replacements**

### Hip Precautions (If Applicable)

Some patients will need to follow hip precautions after surgery. Dr. Burrow/physical therapist will inform you of these precautions if needed.

- **Avoid These Positions**
  - **Do NOT bend your hip past 90 degrees**
    - Avoid leaning forward too far
    - Do not bring your chest toward your knee
  - **Do NOT cross your legs**
    - Keep your legs **separated**
    - Do not cross past the middle of your body
  - **Do NOT rotate your leg inward**
    - Keep your toes pointing **forward or slightly outward**
    - Avoid letting your leg turn inward



## Exercises: First 2 Weeks After Surgery

Perform these exercises until you begin physical therapy or receive further instructions from your physician.

*\*exercise pictures and instructions were chosen from HEP2go.com\**

### Ankle Pumps

- **Frequency:** 10 times per hour while awake
- **How to perform:**
  - Move your foot **up and down at the ankle joint**
- **Why this helps:**
  - Improves blood flow
  - Helps prevent blood clots



### Walking

- Walk using a **walker or crutches**
- **Gradually increase your distance** as tolerated
- **Tips:**
  - Take short, frequent walks
  - Listen to your body—do not overdo it



### Incentive Spirometer (Breathing Exercise)

- **Frequency:** 10 times per hour while awake
- **How to perform:**
  - Exhale normally
  - Place your mouth tightly around the mouthpiece
  - Breathe in **slowly and deeply**
  - Watch the marker rise—try to get it as high as possible
  - Hold your breath for **1 second**, then relax
- **Why this helps:**
  - Keeps your lungs clear
  - Helps prevent pneumonia



*God Bless your recovery! I have already prayed for you during your surgery and will continue to do so. ZPB*